

## Actinin- $\alpha$ 2/3 (Q58) polyclonal antibody

Catalog: BCP00142

Host: Rabbit

Reactivity: Human, Mouse, Rat

### BackGround:

The spectrin gene family encodes a diverse group of cytoskeletal proteins that include spectrins, dystrophins and  $\alpha$ -actinins. There are four tissue-specific  $\alpha$ -actinins, namely  $\alpha$ -actinin-1,  $\alpha$ -actinin-2,  $\alpha$ -actinin-3 and  $\alpha$ -actinin-4, which are localized to muscle and non-muscle cells, including skeletal, cardiac and smooth muscle cells, as well as within the cytoskeleton. Each  $\alpha$ -actinin protein contains one Actin-binding domain, two calponin-homology domains, two EF-hand domains and four spectrin repeats, through which they function as bundling proteins that can cross-link F-Actin, thus anchoring Actin to a variety of intracellular structures. Defects in the gene encoding  $\alpha$ -actinin-4 are the cause of focal segmental glomerulosclerosis 1 (FSGS1), a common renal lesion characterized by decreasing kidney function and, ultimately, renal failure.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 103 kDa

### Swiss-Prot:

P35609/Q08043

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

Actinin- $\alpha$ 2/3 (Q58) polyclonal antibody detects endogenous levels of actinin- $\alpha$ 2 and actinin- $\alpha$ 3 protein.

### DATA:




Western blot (WB) analysis of Actinin- $\alpha$ 2/3 (Q58) polyclonal antibody at 1:500 dilution


Lane1:Hela cell lysate

Lane2:Mouse kidney tissue lysate

Lane3:H9C2 cell lysate




Immunohistochemistry (IHC) analyzes of Actinin- $\alpha$ 2/3 (Q58) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50. showing Nucleus and Cytoplasm staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS1984  
Lot CJ36131

Immunohistochemistry (IHC) analyzes of Actinin- $\alpha$ 2/3 (Q58) pAb in paraffin-embedded human lung carcinoma tissue at 1:50. showing Nucleus and Cytoplasm staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase



Immunohistochemistry (IHC) analyzes of Actinin- $\alpha$ 2/3 (Q58) pAb in paraffin-embedded human kidney carcinoma tissue at 1:50. showing Nucleus and Cytoplasm staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

**Note:**

For research use only, not for use in diagnostic procedure.