

## AKAP 95 (E363) polyclonal antibody

Catalog: BCP00175

Host: Rabbit

Reactivity: Human, Mouse, Rat

### BackGround:

The type II cAMP-protein kinase (PKA) is a multifunctional kinase with a broad range of substrates. Specificity of PKA signaling is thought to be mediated by the compartmentalization of the kinase to specific sites within the cell. To maintain this specific localization, the R subunit (RII) of PKA interacts with specific RII-anchoring proteins. The family of RII-anchoring proteins has been designated A-kinase anchoring proteins (AKAP). AKAP 95, also known as AKAP 8, is a nuclear matrix protein predominantly expressed in liver, heart, pancreas, kidney and skeletal muscle. During mitosis, AKAP 95 is recruited to the chromosomes and plays an essential role in mitotic progression. Characteristic of its family, AKAP 95 participates in PKA signaling through an interaction with the RII regulatory subunit. In addition, AKAP 95 forms a complex with HA95 and HDAC3 and is required for the deacetylation of Histone H3 in mitosis.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 76 kDa

### Swiss-Prot:

O43823

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

AKAP 95 (E363) pAb detects endogenous levels of AKAP 95 (E363) protein.

### DATA:



Western blot (WB) analysis of AKAP 95 (E363) pAb at 1:500 dilution

Lane1:EC9706 whole cell lysate(40ug)

Lane2:HepG2 whole cell lysate(40ug)

Lane3:HEK293T whole cell lysate(40ug)

Lane4:AML-12 whole cell lysate(40ug)

Lane5:H9C2 whole cell lysate(40ug)

### Note:

For research use only, not for use in diagnostic procedure.