

AKT2 (phospho-S474) polyclonal antibody

Catalog: BCP00200 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

AKT, also known as protein kinase B (PKB), is a 57 kDa serine/threonine protein kinase. There are three mammalian isoforms of Akt: AKT1 (PKB alpha), AKT2 (PKB beta) and AKT3 (PKB gamma) with AKT2 and AKT3 being approximately 82% identical with the AKT1 isoform. Each isoform has a pleckstrin homology (PH) domain, a kinase domain and a carboxy terminal regulatory domain. AKT was originally cloned from the retrovirus AKT8, and is a key regulator of many signal transduction pathways. Its tight control over cell proliferation and cell viability are manifold; overexpression or inappropriate activation of AKT has been seen in many types of cancer.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 60 kDa

Swiss-Prot:

P31751

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000 IHC: 1:50~1:200

Storage&Stability:


Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

p-AKT2 (S474) polyclonal antibody detects endogenous levels of AKT2 protein only when phosphorylated at Ser474. This antibody does not cross-react with AKT1 and AKT3 protein when phosphorylated at the corresponding

residues.

DATA:



Western blot (WB) analysis of p-AKT2(S474) pAb at 1:500 dilution Lane1:HEK293T whole cell lysate(40ug)

 $Lane 2: HEK 293T \ treated \ with \ EGF (100 ng/ml, 15 \ minutes) \ whole \ cell \\ ly sate (40 ug)$


Lane3:HEK293T treated with EGF(100ng/ml,30 minutes) whole cell lysate(40ug)

Lane4: The Brain tissue lysate of Rat(40ug)

Lane5: The Brain tissue lysate of Mouse(40ug)

Lane6:C6 whole cell lysate(40ug)

Lane7:AML-12 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-AKT2 (S474) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note

For research use only, not for use in diagnostic procedure.