

## ARNTL polyclonal antibody

Catalog: BCP00232

Host: Rabbit

Reactivity: Human

### BackGround:

Circadian rhythms govern many key physiological processes that fluctuate with a period of approximately 24 hours. These processes include the sleep-wake cycle, glucose, lipid and drug metabolism, heart rate, hormone secretion, renal blood flow, and body temperature, as well as basic cellular processes such as DNA repair and the timing of the cell division cycle. The mammalian circadian system consists of many individual tissue-specific clocks (peripheral clocks) that are controlled by a master circadian pacemaker residing in the suprachiasmatic nuclei (SCN) of the brain. The periodic circadian rhythm is prominently manifested by the light-dark cycle, which is sensed by the visual system and processed by the SCN. The SCN processes the light-dark information and synchronizes peripheral clocks through neural and humoral output signals.

The cellular circadian clockwork consists of interwoven positive and negative regulatory loops, or limbs. The positive limb includes the CLOCK and BMAL1 proteins, two basic helix-loop-helix-PAS containing transcription factors that bind E box enhancer elements and activate transcription of their target genes. CLOCK is a histone acetyltransferase (HAT) protein, which acetylates both histone H3 and H4. BMAL1 binds to CLOCK and enhances its HAT activity. The CLOCK/BMAL1 dimer exhibits a periodic oscillation in both nuclear/cytoplasmic localization and protein levels, both of which are regulated by phosphorylation. CLOCK/BMAL1 target genes include the Cry and Per genes, whose proteins form the negative limb of the circadian clockwork system. CRY and PER proteins (CRY1, CRY2, PER1, PER2 and PER3) form oligomers that also periodically shuttle between the nucleus and cytoplasm. When in the nucleus, CRY/PER proteins inhibit CLOCK/BMAL1-mediated transcriptional activation, thus completing the circadian transcriptional loop. In tissues, roughly six to eight percent of all genes exhibit a circadian expression pattern. This 24-hour periodicity in gene expression results from coordination of the positive and negative regulatory limbs

of the cellular clockwork system, and is fine-tuned by outside signals received from the SCN.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

### Molecular Weight:

~ 85 kDa

### Swiss-Prot:

O00327

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:2000~1:5000

IP 1:50 - 1:100


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

ARNTL polyclonal antibody detects endogenous levels of ARNTL protein.

### DATA:




Western blot (WB) analysis of ARNTL polyclonal antibody at 1:2000 dilution

Lane1:Jurkat whole cell lysate(40ug)

Lane2:A549 whole cell lysate(40ug)

Lane3:Hela whole cell lysate(40ug)



Immunoprecipitation of Jurkat cell lysates using ARNTL pAb (Sepharose Bead Conjugate)#BD0048 (lane 3 and lane 4) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 1 and lane 2). Lane 5 is 30% input. The western blot was probed using ARNTL pAb.

**Note:**

For research use only, not for use in diagnostic procedure.