

BACH1 (L160) polyclonal antibody

Catalog: BCP00261

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

BACH1 and BACH2, heterodimerization partners of MafK, are members of a novel family of BTB/POZ-basic region leucine zipper (bzip) factors. BACH1 and BACH2 have significant similarity to each other in BTB domain and Cap'n' collar-type bZip domain but are otherwise divergent. BACH1 appears ubiquitous, whereas BACH2 is restricted to monocytes, neuronal cells and is abundantly expressed in the early stages of B-cell differentiation. BACH2, a 841 amino acid polypeptide, is an Nrf2-related transcription repressor and a tissue-specific partner of the Maf oncoprotein family.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 90 kDa

Swiss-Prot:

O14867

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1000

IHC: 1:50~1:200


IF: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

BACH1 (L160) polyclonal antibody detects endogenous levels of BACH1 protein.

DATA:

Western blot (WB) analysis of IKBKG pAb at 1:500 dilution

Lane1:The lung tissue lysate of Mouse(40ug)


Lane2:Aml-12 whole cell lysate(40ug)

Lane3:PMVEC whole cell lysate(40ug)

Lane4:HEK293T whole cell lysate(40ug)

Lane5:Hela whole cell lysate (40ug)

Lane6:SHSY5Y whole cell lysate (40ug)



Immunohistochemistry (IHC) analyzes of BACH1 (L160) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50, showing Nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.