# 

## **BMAL2** polyclonal antibody

Catalog: BCP00286

**BackGround:** 

Host: Rabbit

BMAL2, also known as ARNTL2 (aryl hydrocarbon re-

ceptor nuclear translocator-like 2), MOP9, CLIF or

PASD9, is a 636 amino acid protein that localizes to the

nucleus and contains one bHLH (basic helix-loop-helix)

domain, one PAC (PAS-associated C-terminal) domain

and two PAS (PER-ARNT-SIM) domains. Expressed at

high levels in placenta and brain and at lower levels in

liver, thymus, heart, lung and kidney, BMAL2 functions

as a component of the circadian core oscillator, which in-

cludes a variety of proteins that work in tandem to activate the transcription of target genes. More specifically,

BMAL2, when functioning as a component of the core oscillator, binds to the E-box element (3'-CACGTG-5') of

target DNA, thus inducing transcription. Multiple

isoforms of BMAL2 exist due to alternative splicing

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide,

Reactivity: Human

munogen and the purity is > 95% (by SDS-PAGE).

**Applications:** 

WB: 1:2000~1:5000


Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

### **Specificity:**

BMAL2 polyclonal antibody detects endogenous levels of BMAL2 protein.

#### DATA:



Western blot (WB) analysis of BMAL2 polyclonal antibody at 1:2000 dilution

Lane1:PC3 whole cell lysate(40ug)

Lane2:A549 whole cell lysate(40ug)

#### Note:

For research use only, not for use in diagnostic procedure.

### **Molecular Weight:**

events. Product:

~ 70 kDa

**Swiss-Prot:** 

Q8WYA1

**Purification&Purity:** 

50% glycerol, pH7.2.

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-