

Caper (R199) polyclonal antibody

Catalog: BCP00326

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Caper, also known as splicing factor HCC1 or hepatocellular carcinoma protein 1 and RNA binding region containing protein 2 (RNPC2), acts as a transcriptional coactivator for steroid nuclear receptors c-Jun, ER α and ER β . Caper, a nuclear protein with highest concentrations in nuclear speckles, plays a role in the pre-mRNA splicing process. Two isoforms of Caper, HCC1.3 and HCC1.4, co-localize with pre-mRNA splicing factor SC35 and uridine-rich small nuclear RNAs. Caper is a widely expressed protein with highest levels detected in skeletal muscle, lung, brain and pancreas.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 39 kDa

Swiss-Prot:

O75787

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Caper (R199) pAb detects endogenous levels of Caper (R199) protein.

DATA:



Western blot (WB) analysis of Caper (R199) pAb at 1:500 dilution

Lane1:C6 whole cell lysate(40ug)

Lane2:3T3-L1 whole cell lysate(40ug)

Lane3:HepG2 whole cell lysate(40ug)

Lane4:U-87MG whole cell lysate(40ug)

Lane5:HEK293T whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.