

CD105 polyclonal antibody

Catalog: BCP00366

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

Vascular endothelium glycoprotein that plays an important role in the regulation of angiogenesis. Required for normal structure and integrity of adult ture. Regulates the migration of vascular endothelial cells. Required for normal extraembryonic angiogenesis and for embryonic heart development (By similarity). May regulate endothelial cell shape changes in response to blood flow, which drive vascular remodeling and establishment of normal vascular morphology during angiogenesis (By similarity). May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors. Acts as TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade that ultimately leads to the activation of SMAD transcription tors. Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGFB1 signaling through SMAD3.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~90 kDa

Swiss-Prot:

P17813

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000~1:2000

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Specificity:

CD105 polyclonal antibody detects endogenous levels of CD105 protein.


DATA:



Western blot (WB) analysis of CD105 polyclonal antibody at 1:1000 dilution

Lane 1: MCF-7 cell membrane lysate (20ug)

Lane 2: A549 cell membrane lysate (30ug)



Western blot (WB) analysis of CD105 polyclonal antibody at 1:1000 dilution

Lane 1: A549 whole cell lysate (40ug)

Lane 2: CT-26 whole cell lysate (40ug)

Note:

For research use only, not for use in diagnostic procedure.