

# CD114/CSF3R polyclonal antibody

Catalog: BCP00372 Host: Rabbit Reactivity: Human, Mouse

#### **BackGround:**

Granulocyte colony-stimulating factor (G-CSF-R) is a transmembrane protein comprised of an immunoglobulin-like (Ig-like) domain, a cytokine receptor-homologous (CRH) domain, and three fibronectin type III (FN III) domains. G-CSF-R is expressed on all granulocytic lineages, including progenitor cells, and has been detected on monocytes, T and B lymphocytes, as well as non-hematopoietic tissues including cardiomyocytes and neural stem cells. The primary ligand for G-CSF-R is the cytokine granulocyte colony-stimulating factor (G-CSF). G-CSF-R has no intrinsic tyrosine kinase activity; ligand binding induces conformational changes in the receptor, leading to activation of the Jak/Stat, PI3K/Akt, and MAPK pathways. G-CSF induces differentiation and proliferation of myeloid progenitor cells into neutrophils. Multiple diseases have been associated with mutations of the G-CSF-R gene, CSF3R, including severe congenital neutropenia (SCN), chronic neutrophilic leukemia (CNL), and atypical chronic myeloid leukemia (aCML).

## **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

# **Molecular Weight:**

~ 105 kDa

## **Swiss-Prot:**

Q99062

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum

by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

## **Applications:**

WB: 1:1000~1:2000


# Storage&Stability:

Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

## **Specificity:**

CD114/CSF3R polyclonal antibody detects endogenous levels of CD114/CSF3R protein.

#### **DATA:**



Western blot (WB) analysis of CD114/CSF3R polyclonal antibody at 1:1000 dilution

Lane1:A549 whole cell lysate(30ug)

Lane2:SP2/0 whole cell lysate(30ug)

Lane3:The Brain tissue lysate of Mouse(30ug)

#### Note:

For research use only, not for use in diagnostic procedure.