

COUP-TFI polyclonal antibody

Catalog: BCP00556 Host: Rabbit Reactivity: Pig

BackGround:

Chicken ovalbumin upstream promoter transcription factor (COUP-TF) belongs to the NR2 subfamily of the nuclear hormone receptor family. COUP-TFI and COUP-TFII are two of the well-characterized members in the NR2 subfamily. These two members are highly conserved in their two zinc-finger DNA binding domains (DBD) and the ligand binding domain (LBD), and function as repressors or activators of downstream target genes to regulate different biological processes. COUP-TFI and II bind to 5'-AGGTCA-3' motif palindromes, either directly or indirectly, through heterodimer formation with other proteins (e.g. RXRs) to regulate downstream target gene expression. COUP-TFI is involved in neuronal development, tissue patterning, and differentiation. COUP-TFII has been shown to be involved in angiogenesis, glucose homeostasis, and mesenchymal cell commitment.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 52 kDa

Swiss-Prot:

P24468

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:5000~1:10000


Storage&Stability:

Store at $4 \,^{\circ}$ C short term. Aliquot and store at $-20 \,^{\circ}$ C long term. Avoid freeze-thaw cycles.

Specificity:

COUP-TFI polyclonal antibody detects endogenous levels of COUP-TFI protein.

DATA:



Western blot (WB) analysis of COUP-TFI polyclonal antibody at

1:10000 dilution

Lane1:The Heart tissue lysate of Pig(40ug)

Lane2: The Brain tissue lysate of Pig(40ug)

Note:

For research use only, not for use in diagnostic procedure.