

EEF2 (A50) polyclonal antibody

Catalog: BCP00684

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Two elongation factors (EF) EF-Tu and EF-2 participate in the elongation phase during protein biosynthesis on the ribosome and their functional cycles depend on GTP binding and its hydrolysis. EF-Tu (also designated mitochondrial precursor p43) and EF-2 are multidomain GTPases with essential functions in translation, and they both bind to the same site on the ribosome where their low intrinsic GTPase activities are strongly stimulated. EF-Tu plays a central role in the fast and accurate delivery of aminoacyl-tRNAs to the translating ribosome. In addition, EF-Tu protects the aminoester bond against hydrolysis until a correct match between the codon on mRNA and the anticodon on tRNA can be achieved. EF-2 supports the translocation of tRNAs and of mRNAs on the ribosome so that a new codon can be exposed for decoding.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 95 kDa

Swiss-Prot:

P13639

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

eEF2 (A50) polyclonal antibody detects endogenous levels of eEF2 protein.

DATA:




Western blot (WB) analysis of eEF2 (A50) polyclonal antibody at 1:500 dilution

Lane1:HepG2 cell lysate

Lane2:Mouse kidney tissue lysate

Lane3:Rat kidney tissue lysate



Immunohistochemistry (IHC) analyzes of eEF2 (A50) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.