

EGFR (phospho-Y992) polyclonal antibody

Catalog: BCP00695

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Epidermal growth factor mediates its effects on cell growth through its interaction with a cell surface glycoprotein designated the EGF receptor. Binding of EGF or TGF α to the EGF receptor activates tyrosine-specific protein kinase activity intrinsic to the EGF receptor. The carboxy-terminal tyrosine residues on EGFR, Tyr 1068 and Tyr 1173, are the major sites of autophosphorylation, which occurs as a result of EGF binding. Phosphorylation of Tyr 992, Tyr 1068 and Tyr 1086 is required for conformational change in the C-terminal tail of the EGF receptor regulates Actin filament accumulation at the plasma membrane and Cdc42 stimulates formation of filopodia.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 175 kDa

Swiss-Prot:

P00533

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000 IHC: 1:50~1:200 IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-EGFR (Y992) polyclonal antibody detects endogenous levels of EGFR protein only when phosphorylated at Tyr992 (removal of the signal peptide).


DATA:




Western blot (WB) analysis of p-EGFR (Y992) pAb at 1:500 dilution

Lane1:HCC827 whole cell lysate(20ug)

Lane2:A549 whole cell lysate(40ug)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using p-EGFR (Y992) pAb. The picture on the right is treated with the synthesized peptide.



Immunofluorescence analysis of HeLa cells, using p-EGFR (Y992) pAb. The picture on the right is treated with the synthesized peptide.

Note:

For research use only, not for use in diagnostic procedure.