

eIF3ε (V116) polyclonal antibody

Catalog: BCP00704

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Translation initiation in eukaryotes necessitates the assembly of an 80S ribosomal complex containing methionyl initiator tRNA (Met-tRNAⁱMet), which is base paired at the initiation codon (AUG, GUG) in eligible transcripts. Eukaryotic initiation factors (eIFs) are utilized in a sequence of reactions that leads to 80S ribosomal assembly and initiation of translation. Eukaryotic initiation factor 3 (eIF3) is the largest family of eIFs and consists of at least 12 unique subunits in mammals. eIFε, also known as eIF p47, binds to the 40S ribosome and promotes the binding of methionyl-tRNAⁱ and mRNA and associates with the complex p170-eIF3.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 47 kDa

Swiss-Prot:

O00303

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


IHC: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

eIF3ε (V116) polyclonal antibody detects endogenous levels of eIF3ε protein.

DATA:


Western blot (WB) analysis of eIF3ε (V116) pAb at 1:1000 dilution

Lane1: The Testis tissue lysate of Mouse(40ug)

Lane2: The Testis tissue lysate of Rat(40ug)

Lane3: Hela whole cell lysate(40ug)

Lane4: Panc1 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of eIF3ε (V116) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50.

Note:

For research use only, not for use in diagnostic procedure.