

eIF3 ζ (R135) polyclonal antibody

Catalog: BCP00705

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Translation initiation in eukaryotes necessitates the assembly of an 80S ribosomal complex containing methionyl initiator tRNA (Met-tRNAⁱMet), which is base paired at the initiation codon (AUG, GUG) in eligible transcripts. Eukaryotic initiation factors (eIFs) are utilized in a sequence of reactions that leads to 80S ribosomal assembly and initiation of translation. Eukaryotic initiation factor 3 (eIF3) is the largest family of eIFs and consists of at least 12 unique subunits in mammals. eIF ϵ , also known as eIF p47, binds to the 40S ribosome and promotes the binding of methionyl-tRNAⁱ and mRNA and associates with the complex p170-eIF3.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 64 kDa

Swiss-Prot:

O15371

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

eIF3 ζ (R135) polyclonal antibody detects endogenous levels of eIF3 ζ protein.

DATA:




Western blot (WB) analysis of eIF3 ζ (R135) pAb at 1:1000 dilution

Lane1:The Brain tissue lysate of Mouse(40ug)

Lane2:The Brain tissue lysate of Rat(20ug)

Lane3:HEK293T whole cell lysate(10ug)

Lane4:MCF-7 whole cell lysate(20ug)



Immunohistochemistry (IHC) analyzes of eIF3 ζ (R135) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.