

FMIP (E602) polyclonal antibody

Catalog: BCP00780

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

Although the macrophage colony stimulating factor (M-CSF) and its receptor, c-Fms, are involved in the survival and proliferation of hematopoietic cells, little is known about the signalling events leading to differentiation into mature blood cells. A 78 kDa Fms-interacting protein, FMIP, transiently binds to M-CSF-activated Fms-molecules. This binding results in a rapid phosphorylation of FMIP within its Fms-binding domain, thereby dissociating Fms and FMIP. Endogenous levels of FMIP may form a threshold that decide whether bipotential progenitor cells differentiate into macrophages or granulocytes. Myeloid progenitor cells express low levels of endogenous FMIP and, upon M-CSF specific signalling, are differentiated into macrophages. Overexpression of FMIP may saturate Fms, which results in predominant cytoplasmic expression of FMIP and favors granulocyte differentiation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 78 kDa

Swiss-Prot:

Q13769

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

FMIP (E602) polyclonal antibody detects endogenous levels of FMIP protein.

DATA:




Western blot (WB) analysis of FMIP (E602) polyclonal antibody at 1:500 dilution

Lane1:Hela whole cell lysate

Lane2:NIH-3T3 whole cell lysate

Lane3:PC12 whole cell lysate



Immunohistochemistry (IHC) analyzes of FMIP (E602) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.