

Gads (N158) polyclonal antibody

Catalog: BCP00802

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

The Src homology 3 (SH3) region is a small protein domain of approximately 60 amino acids present in a large group of proteins. In general, it exists in association with catalytic domains, as in the nonreceptor protein-tyrosine kinases and phospholipase C- γ , within structural proteins, such as spectrin or myosin, and in small adapter proteins, such as Crk and GRB2. SH3 domains are often accompanied by SH2 domains of 100 amino acids that bind to tyrosine-phosphorylated regions of target proteins, frequently linking activated growth factors to putative signal transduction proteins. Deletion or mutation of SH3 domains generally activates the transforming potential of nonreceptor tyrosine kinases, suggesting that SH3 mediates negative regulation of an intrinsic transforming activity. Gads is an adapter proteins that contains both SH2 and SH3 domains. Gads binds to tyrosine-phosphorylated proteins, such as Shc, and functions to couple these proteins to downstream effectors.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 38 kDa

Swiss-Prot:

O75791

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Gads (N158) polyclonal antibody detects endogenous levels of Gads protein.

DATA:



Western blot (WB) analysis of Gads (N158) pAb at 1:1000 dilution

Lane1:CT26 whole cell lysate(40ug)

Lane2:PC12 whole cell lysate(40ug)

Lane3:SGC7901 whole cell lysate(40ug)

Lane4:H1792 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.