

GATA-6 (R493) polyclonal antibody

Catalog: BCP00809

Host: Rabbit

Reactivity: Mouse,Rat

BackGround:

Members of the GATA family share a conserved zinc finger DNA-binding domain and are capable of binding the WGATAR consensus sequence. GATA-1 is erythroid-specific and is responsible for the regulated transcription of erythroid genes. It is an essential component in the generation of the erythroid lineage. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as erythroid cells. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1. GATA-3 is highly expressed in T-lymphocytes. GATA-4, GATA-5 and GATA-6 comprise a subfamily of transcription factors. GATA-4 and GATA-6 are found in heart, pancreas and ovary; lung and liver tissues exhibit GATA-6, but not GATA-4, expression.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 55 kDa

Swiss-Prot:

Q92908

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum

by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

GATA-6 (R493) polyclonal antibody detects endogenous levels of GATA-6 protein.

DATA:



Western blot (WB) analysis of GATA-6 (R493) pAb at 1:500 dilution

Lane1: The Heart tissue lysate of Mouse(40ug)

Lane2: The Heart tissue lysate of Rat(40ug)

Note:

For research use only, not for use in diagnostic procedure.