

GCSc- γ (E315) polyclonal antibody

Catalog: BCP00812

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

The GCLC gene consists of 16 exons and encodes the 636 amino acid protein γ -GCSc (γ -glutamylcysteine synthetase heavy subunit), also designated γ -L-glutamate-L-cysteine ligase catalytic subunit (GLCLC). γ -GCSc is expressed in hemocytes, brain, liver and kidney. γ -GCSc associates with a regulatory or modifier subunit, γ -GCScm (γ -glutamylcysteine synthetase light subunit), to form a heterodimer, γ -GCS. γ -GCS is the first enzyme involved and the rate determining step in glutathione biosynthesis. Oxidants, cadmium and methyl mercury upregulate the transcription of γ -GCS. H₂O₂ regulation depends on the Yap1 protein and the presence of glutamate, glutamine and lysine. Cadmium regulates transcription through proteins Met-4, Met-31 and Met-32. Cbf1, a DNA binding protein, inhibits transcription of γ -GCS.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 70 kDa

Swiss-Prot:

P48506

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

GCSc- γ (E315) polyclonal antibody detects endogenous levels of GCSc- γ protein.

DATA:



Western blot (WB) analysis of GCSc- γ (E315) polyclonal antibody at 1:500 dilution

Lane1:HepG2 whole cell lysate(40ug)

Lane2:NIH-3T3 whole cell lysate(40ug)

Lane3:PC12 whole cell lysate(40ug)

Lane4:A549 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.