

HDAC3 (Y414) polyclonal antibody

Catalog: BCP00859 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, PCAF (p300/CBP-associated factor), p300/CBP and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated RPD3) and HDAC3, all of which are related to the yeast transcriptional factor Rpd3p, have been identified as histone deacetylases.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 49 kDa

Swiss-Prot:

O15379

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000 IHC: 1:50~1:200 IF: 1:50~1:200


Storage&Stability:

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

HDAC3 (Y414) polyclonal antibody detects endogenous levels of HDAC3 protein.

DATA:




Western blot (WB) analysis of HDAC3 (Y414) polyclonal antibody at 1:500 dilution

Lane1:COS-7 whole cell lysate(40ug)

Lane2:Hela whole cell lysate(40ug)

Lane3:MEF whole cell lysate(40ug)

Lane4:C6 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of HDAC3 (Y414) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.