

HNF4 α polyclonal antibody

Catalog: BCP00890

Host: Rabbit

Reactivity: Human

BackGround:

Hepatocyte nuclear factor 4 α (HNF4 α) is a transcription factor that belongs to the steroid hormone receptor super-family and is enriched in liver. HNF4 α , in association with PGC-1 α , activates gluconeogenic genes such as phosphoenolpyruvate carboxykinase and glucose-6-phosphatase genes in fasted livers. Conditional knockout of the HNF4 α gene in the mouse liver destroys lipid homeostasis and leads to lipid accumulation in the liver and a reduction of serum cholesterol and triglyceride levels. Mutations in HNF4 α have been linked to maturity-onset diabetes of the young (MODY).

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 58 kDa

Swiss-Prot:

P41235

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000~1:2000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

HNF4 α polyclonal antibody detects endogenous levels of HNF4 α protein.


DATA:



Western blot (WB) analysis of HNF4 α polyclonal antibody at 1:2000 dilution

Lane1:HepG2 whole cell lysate(40ug)

Lane2:A549 whole cell lysate(40ug)



Immunohistochemistry of paraffin-embedded Human Brain using HNF4 α antibody at dilution of 1:50.

Note:

For research use only, not for use in diagnostic procedure.