

## IKK $\alpha$ (M17) polyclonal antibody

Catalog: BCP00940

Host: Rabbit

Reactivity: Human, Mouse, Rat

### BackGround:

The NF- $\kappa$ B/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory I $\kappa$ B proteins. Most agents that activate NF- $\kappa$ B do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of I $\kappa$ B. The key regulatory step in this pathway involves activation of a high molecular weight I $\kappa$ B kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits. IKK $\alpha$  and IKK $\beta$  serve as the catalytic subunits of the kinase and IKK $\gamma$  serves as the regulatory subunit. Activation of IKK depends upon phosphorylation at Ser177 and Ser181 in the activation loop of IKK $\beta$  (Ser176 and Ser180 in IKK $\alpha$ ), which causes conformational changes, resulting in kinase activation.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 85 kDa

### Swiss-Prot:

O15111

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


### Storage&Stability:

Store at 4  $^{\circ}$ C short term. Aliquot and store at -20  $^{\circ}$ C long term. Avoid freeze-thaw cycles.

### Specificity:

IKK- $\alpha$  (M17) polyclonal antibody detects endogenous levels of IKK- $\alpha$  protein.

### DATA:



Western blot (WB) analysis of IKK $\alpha$  (M17) pAb at 1:500 dilution

Lane1:3T3-L1 whole cell lysate(40ug)


Lane2:C6 whole cell lysate(40ug)

Lane3:HEK293T whole cell lysate(40ug)

Lane4:L02 whole cell lysate(40ug)

Lane5:HepG2 whole cell lysate(40ug)

Lane6:HCT116 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of IKK $\alpha$  (M17) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

### Note:

For research use only, not for use in diagnostic procedure.