

IKK γ (H81) polyclonal antibody

Catalog: BCP00944

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

Activation of NF κ B requires that I κ B be phosphorylated on specific serine residues, which results in targeted degradation of I κ B. I κ B kinase α (IKK α), previously designated CHUK, interacts with I κ B- α and specifically phosphorylates I κ B α on Serine 32 and 36, the sites that trigger its degradation. IKK α appears to be critical for NF κ B activation in response to proinflammatory cytokines. Phosphorylation of I κ B by IKK α is stimulated by the NF κ B inducing kinase (NIK), which itself is a central regulator for NF κ B activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKK α , IKK β and IKK γ (also designated NEMO), and each appear to make essential contributions to I κ B phosphorylation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 42 kDa

Swiss-Prot:

Q9Y6K9

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 $^{\circ}$ C short term. Aliquot and store at -20 $^{\circ}$ C long term. Avoid freeze-thaw cycles.

Specificity:

IKK γ (H81) polyclonal antibody detects endogenous levels of IKK γ protein.

DATA:



Western blot (WB) analysis of IKK γ pAb at 1:500 dilution


Lane1:MEF whole cell lysate(40ug)

Lane2:H9C2 whole cell lysate(40ug)

Lane3:H1792 whole cell lysate(40ug)

Lane4:HEK293T whole cell lysate(40ug)

Lane5:Panc1 whole cell lysate (40ug)



Immunohistochemistry (IHC) analyzes of IKK γ (H81) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.