

IRAK1 (T381) polyclonal antibody

Catalog: BCP00975

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Three structurally related ligands for IL-1Rs have been described. These include two agonists, IL-1 α and IL-1 β , and a specific receptor antagonist, IL-1Ra. Two distinct receptors designated IL-1RI and IL-1RII have been identified, each of which belong to the Ig superfamily. The preponderance of evidence suggests IL-1RI to be the functional IL-1 receptor. Binding of IL-1 to its cognate receptor results in the activation of the NF κ B signaling pathway. The IL-1-dependent kinase termed IRAK (for IL-1 receptor-associated kinase) co-immunoprecipitates with activated IL-1RI and has been implicated as an upstream mediator of NF κ B activation. Additional support for this assertion comes from the fact that a related Drosophila protein, Pelle, is a known upstream activator of Dorsal, the Drosophila homolog of NF κ B.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 76 kDa

Swiss-Prot:

P51617

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

IRAK1 (T381) polyclonal antibody detects endogenous levels of IRAK1 protein.

DATA:



Western blot (WB) analysis of IRAK1 (T381) polyclonal antibody at 1:500 dilution

Lane1:L02 whole cell lysate(40ug)

Lane2:THP-1 whole cell lysate(40ug)

Lane3:HCT116 whole cell lysate(40ug)

Lane4:PC12 whole cell lysate(40ug)

Lane5:CT26 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.