

IRS-1 (phospho-S323) polyclonal antibody

Catalog: BCP00984

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

IRS-1, a major substrate of the insulin receptor, is phosphorylated in response to stimulation of cells by insulin, insulin-like growth factor 1 (IGF-1) and interleukin 4 (IL-4). IRS-1 is phosphorylated on serine, threonine and tyrosine residues in a variety of tissues. An insulin-sensitive serine/threonine kinase casein kinase II mediates a portion of the insulin-stimulated serine/threonine phosphorylation of overexpressed IRS-1 in vivo. Thr 502 is identified as the major casein kinase II-catalyzed phosphorylation site in rat IRS-1, and Ser 99 is an additional phosphorylation site catalyzed by casein kinase II. Thus, casein kinase II-catalyzed phosphorylation of IRS-1 may be a component of the intracellular insulin signaling cascade. IRS-1 contains three putative binding sites for 14-3-3 (Ser 270, Ser 374 and Ser 641) and the motif around Ser 270 is located in the phosphotyrosine binding domain of IRS-1, which is responsible for the interaction with the insulin receptor.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 132, 180 kDa

Swiss-Prot:

P35568

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-IRS-1 (S323) polyclonal antibody detects endogenous levels of IRS-1 only when phosphorylated at Ser323.

DATA:



Western blot (WB) analysis of p-IRS-1 (S323) polyclonal antibody at 1:500 dilution


Lane1:L02 whole cell lysate(40ug)

Lane2:L02 treated with PBS(1×PBS,PH7.4) for 1 heure whole cell lysate

Lane3:L02 treated with PBS(1×PBS,PH7.4) for 1 heure then treated with DMEM(10%FBS) for 5min whole cell lysate

Lane4:L02 treated with PBS(1×PBS,PH7.4) for 1 heure then treated with DMEM(10%FBS) for 10min whole cell lysate

Lane5:L02 treated with PBS(1×PBS,PH7.4) for 1 heure then treated with DMEM(10%FBS) for 15min whole cell lysate



Immunohistochemistry (IHC) analyzes of p-IRS-1 (S323) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.