

JNK3 (K391) polyclonal antibody

Catalog: BCP01001 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

MAPK10 (JNK3) is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This protein is a neuron-specific form of c-Jun N-terminal kinases (JNKs). Through its phosphorylation and nuclear localization, this kinase plays regulatory roles in the signaling pathways of neuronal apoptosis. Beta-arrestin 2, a receptor-regulated MAP kinase scaffold protein, is found to interact with and stimulate the phosphorylation of this kinase by MAP kinase kinase 4 (MKK4). Cyclin-dependent kianse 5 (CDK5) can phosphorylate and inhibit the activity of this kinase, which may be important in preventing neuronal apoptosis. Four alternatively spliced transcript variants encoding distinct isoforms have been reported.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 53 kDa

Swiss-Prot:

P53779

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000 IHC: 1:50~1:200 IF: 1:50~1:200


Storage&Stability:

Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

JNK3 (K391) polyclonal antibody detects endogenous levels of JNK3 protein.

DATA:




Western blot (WB) analysis of JNK3 (K391) pAb at 1:1000 dilution

Lane1:C6 whole cell lysate(40ug)

Lane2:CT26 whole cell lysate(40ug)

Lane3:U-87MG whole cell lysate(40ug)

Lane4:MCF-7 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of JNK3 (K391) pAb in paraffin-embedded human brain tissue.

Note:

For research use only, not for use in diagnostic procedure.