

MGMT (C5) polyclonal antibody

Catalog: BCP01096

Host: Rabbit

Reactivity: Human

BackGround:

MGMT (O6-methylguanine-DNA methyltransferase) is transcriptionally activated in response to DNA damage and functions to repair mutagenic and cytotoxic O6-alkylguanine lesions caused by carcinogens or cytostatic drugs. MGMT induction by ionising radiation does not occur in p53-deficient mice, suggesting that MGMT induction may require p53. Similarly, MGMT mRNA and protein were shown to be inducible by ionising radiation only in cell lines that express functional p53, and not in cell lines that do not express wild type p53. In contrast, in a study of oral cancer cell lines, high MGMT activity was associated with the presence of mutant p53. Similarly, MGMT activity was significantly lower in ovarian tumors with wild-type p53 than in tumors with mutant p53, supporting the view that wild type p53 down-regulates the basal MGMT promoter.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 22 kDa

Swiss-Prot:

P16455

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

MGMT (C5) polyclonal antibody detects endogenous levels of MGMT protein.

DATA:



Western blot (WB) analysis of MGMT (C5) pAb at 1:500 dilution


Lane1:L02 whole cell lysate(40ug)

Lane2:A549 whole cell lysate(40ug)

Lane3:PC3 whole cell lysate(40ug)

Lane4:CT26 whole cell lysate(40ug)

Lane5:PC12 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of MGMT (C5) pAb in paraffin-embedded human colon carcinoma tissue at 1:50, showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.