

Mob1A/B (D127) polyclonal antibody

Catalog: BCP01109

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

MOB1 was first identified in yeast as a protein that binds to Mps with essential roles in the completion of mitosis and the maintenance of ploidy. Its Drosophila and mammalian homologs, Mats and MOB1, respectively, are involved in the Hippo signaling tumor suppressor pathway, which plays a critical role in organ size regulation and which has been implicated in cancer development. There are two MOB1 proteins in humans, MOB1 α and MOB1 β , that are encoded by two different genes but which have greater than 95% amino acid sequence identity. Both forms bind to members of the nuclear Dbf2-related (NDR) kinases, such as LATS1/2 and NDR1/2, thereby stimulating kinase activity. This binding is promoted by the phosphorylation of MOB1 at several threonine residues by MST1 and/or MST2.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 25 kDa

Swiss-Prot:

Q7L9L4

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Mob1A/B (D127) polyclonal antibody detects endogenous levels of Mob1A/B protein.

DATA:



Western blot (WB) analysis of Mob1A/B (D127) polyclonal antibody at 1:500 dilution

Lane1:HeLa whole cell lysate(40ug)

Lane2:H1792 whole cell lysate(20ug)

Lane3:HCT116 whole cell lysate(40ug)

Lane4:CT26 whole cell lysate(40ug)

Lane5:PMVEC whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.