

p38 MAPK (Ab-180) polyclonal antibody

Catalog: BCP01237

Host: Rabbit

Reactivity: Human,Rat,Mouse

BackGround:

p38 MAP kinase (MAPK), also called RK or CSBP, is the mammalian orthologue of the yeast HOG kinase that participates in a signaling cascade controlling cellular responses to cytokines and stress. Four isoforms of p38 MAPK, p38 α , β , γ (also known as Erk6 or SAPK3), and δ (also known as SAPK4) have been identified. Similar to the SAPK/JNK pathway, p38 MAPK is activated by a variety of cellular stresses including osmotic shock, inflammatory cytokines, lipopolysaccharide (LPS), UV light, and growth factors. MKK3, MKK6, and SEK activate p38 MAPK by phosphorylation at Thr180 and Tyr182. Activated p38 MAPK has been shown to phosphorylate and activate MAPKAP kinase 2 and to phosphorylate the transcription factors ATF-2, Max, and MEF2.

SB203580

(4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-imidazole) is a selective inhibitor of p38 MAPK. This compound inhibits the activation of MAPKAPK-2 by p38 MAPK and subsequent phosphorylation of HSP27. SB203580 inhibits p38 MAPK catalytic activity by binding to the ATP-binding pocket, but does not inhibit phosphorylation of p38 MAPK by upstream kinases.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 41 kDa

Swiss-Prot:

Q16539

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95%

Applications:

WB: 1:5000~1:10000

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p38 MAPK (Ab-180) polyclonal antibody detects endogenous levels of p38 MAPK protein.

DATA:



Western blot (WB) analysis of p38 MAPK (Ab-180) polyclonal antibody at 1:5000 dilution

Lane1:C6 whole cell lysate(40ug)


Lane2:AML-12 whole cell lysate(40ug)

Lane3:The Heart tissue lysate of Rat(40ug)

Lane4:H1792 whole cell lysate(40ug)

Lane5:MCF-7 whole cell lysate(40ug)

Lane6:Hela whole cell lysate(40ug)



Immunofluorescence analysis of PC3 cells using p38 MAPK antibody at dilution of 1:50.

Note:

For research use only, not for use in diagnostic procedure.