

# PFKFB1/4 (E349) polyclonal antibody

Catalog: BCP01294 Host: Rabbit Reactivity: Human, Mouse, Rat

#### **BackGround:**

PFK-1 undergoes activation in the presence of elevated AMP. The most potent activator is fructose-2,6-bisphosphate, which is produced by PFK-2 from the same substrate, fructose 6-phosphate. PFK-2 is bifunctional and a key regulator for PFK-1. PFK-2 catalyzes the synthesis of fructose-2,6-bisphosphate, and contains fructose-2,6-biphosphatase activity that catalyzes the degradation of fructose-2,6-bisphosphate. PFK-2 is dimeric and isoenzymes include PFK-2 liver (PFKFB1, PFRX), PFK-2 cardiac (PFKFB2), PFK-2 placental (PFKFB3, inducible PFK-2) and PFK-2 testis (PFKFB4).

#### **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

# **Molecular Weight:**

~ 54 kDa

# **Swiss-Prot:**

P16118/Q16877

# **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

# **Applications:**

WB: 1:500~1:1000


# **Storage&Stability:**

Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

# **Specificity:**

PFKFB1/4 (E349) polyclonal antibody detects endogenous levels of PFKFB1/4 protein.

# **DATA:**



Western blot (WB) analysis of PFKFB1/4 (E349) pAb at 1:1000 dilution

Lane1:K562 whole cell lysate(40ug)

Lane2:A549 whole cell lysate(40ug)

Lane3:PC12 whole cell lysate(40ug)

Lane4:AML-12 whole cell lysate(40ug)

### Note:

For research use only, not for use in diagnostic procedure.