

PGD synthase (M115) polyclonal antibody

Catalog: BCP01296

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

Glycogen is a polysaccharide of glucose and serves as an energy storage in mammalian muscle and liver (1). Glycogen synthase catalyzes the rate-limiting step of glycogen biosynthesis and has two major isoforms in mammals -- muscle isoform (GYS1) and liver isoform (GYS2) respectively. Glycogen synthase kinase-3 α (GSK-3 α) and glycogen synthase kinase-3 β (GSK-3 β) phosphorylate glycogen synthase at multiple sites in its C-terminus (Ser641, Ser645, Ser649 and Ser653) inhibiting its activity. Hypoxia alters glycogen metabolism including temporal changes of GYS1 expression and phosphorylation in cancer cells, suggesting the role of metabolic reprogramming of glycogen metabolism in cancer growth.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 23 kDa

Swiss-Prot:

O60760

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PGD synthase (M115) polyclonal antibody endogenous levels of PGD synthase protein.

DATA:



Western blot (WB) analysis of PGD synthase (M115) polyclonal antibody at 1:500 dilution

Lane1:SK-OVCAR3 whole cell lysate(40ug)

Lane2:U-87MG whole cell lysate(40ug)

Lane3:MCF-7 whole cell lysate(40ug)

Lane4:HepG2 whole cell lysate(40ug)

Lane5:C6 whole cell lysate(40ug)

Lane6:BV2 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.