

PI3K p85 α (Tyr607) polyclonal antibody

Catalog: BCP01304

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

The enzyme phosphatidylinositol 3 kinase (PI3 kinase) is a lipid kinase that generates phosphatidylinositol 3, 4, 5-triphosphate in response to receptor activation in many signal transduction pathways. Class IA PI3Ks exist as a heterodimer of a catalytic 110 kDa (p110) and a regulatory p85 subunit (e.g. p85 alpha). p85 alpha is an adaptor molecule that regulates the activity of the catalytic p110 subunit by binding to phosphorylated receptor tyrosine kinases (RTKs) through its SH2 domain and mediating the interaction between p110 and the plasma membrane. p85 alpha is necessary for insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 52, 85 kDa

Swiss-Prot:

P27986/Q92569

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000

IF: 1:50~1:200

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PI3K p85 α /p55 γ (Tyr607) polyclonal antibody detects

endogenous levels of PI3-kinase p85 α protein. This antibody also detects PI3K p55 γ protein.

DATA:



Western blot (WB) analysis of PI3K p85 α (Tyr607) pAb at 1:2000 dilution


Lane1:C6 whole cell lysate(40ug)

Lane2:CT-26 whole cell lysate(40ug)

Lane3:Hela whole cell lysate(40ug)

Lane4:Jurkat whole cell lysate(40ug)

Lane5:MCF-7 whole cell lysate(40ug)



Immunofluorescence analysis of BV2 cells using PI3K p85 α /p55 γ antibody at dilution of 1:50.

Note:

For research use only, not for use in diagnostic procedure.