

## PIAS3 (F25) polyclonal antibody

Catalog: BCP01307

Host: Rabbit

Reactivity: Human,Rat,Mouse

### BackGround:

The IL-6 type family of cytokines, which includes IL-6 as well as a number of similar cytokines and growth factors, plays a significant role in regulating gene activation, proliferation and differentiation. Transcription factors of the Stat family are known to be involved in this signal transduction pathway, undergoing phosphorylation, dimerization, and translocation to the nucleus upon activation. PIAS 1, for protein inhibitor of activated Stat1 (also designated Gu/RNA helicase II binding protein), binds specifically to Stat1, blocking Stat1 DNA-binding activity and inhibiting Stat1-mediated gene activation. PIAS 1 also binds to the Gu/RNA helicase II enzyme, leading to the proteolytic cleavage of Gu/RH-II. PIAS 3 similarly binds specifically to Stat3, blocking Stat3 DNA-binding activity and inhibiting Stat3-mediated gene activation.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 75 kDa

### Swiss-Prot:

Q9Y6X2

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:5000~1:10000

IF: 1:50~1:200


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

PIAS3 (F25) polyclonal antibody detects endogenous levels of PIAS3 protein.

### DATA:



Western blot (WB) analysis of PIAS3 (F25) polyclonal antibody at 1:5000 dilution


Lane1:A549 whole cell lysate(40ug)

Lane2:PC3 whole cell lysate(40ug)

Lane3:HeK293T whole cell lysate(40ug)

Lane4:H9C2 whole cell lysate(40ug)

Lane5:MEF whole cell lysate(40ug)



Immunofluorescence analysis of HeLa cells using PIAS3 antibody at dilution of 1:50.

### Note:

For research use only, not for use in diagnostic procedure.