

PKA α / β cat (V191) polyclonal antibody

Catalog: BCP01322

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The second messenger cyclic AMP (cAMP) mediates diverse cellular responses to external signals such as proliferation, ion transport, regulation of metabolism and gene transcription by activation of the cAMP-dependent protein kinase (cAPK or PKA). Activation of PKA occurs when cAMP binds to the two regulatory subunits of the tetrameric PKA holoenzyme resulting in release of active catalytic subunits. Three catalytic (C) subunits have been identified, designated C α , C β and C γ , that each represent specific gene products. C α and C β are closely related (93% amino acid sequence similarity), whereas C γ displays 83% and 79% similarity to C α and C β , respectively.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 40, 42 kDa

Swiss-Prot:

P17612/P22694

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


IF: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PKA α / β cat (V191) polyclonal antibody detects endogenous levels of PKA α / β protein.

DATA:


Western blot (WB) analysis of PKA α / β cat (V191) polyclonal antibody at 1:500 dilution

Lane1:HEK293T whole cell lysate(40ug)

Lane2:MCF-7 whole cell lysate(40ug)

Lane3:The Testis tissue lysate of Rat(40ug)

Lane4:The Testis tissue lysate of Mouse(40ug)



Immunohistochemistry (IHC) analyzes of PKA α / β cat (V191) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.