# PKC δ (phospho-S645) polyclonal antibody

Catalog: BCP01329

Host:

Rabbit

Reactivity: Human

# **BackGround:**

Protein Kinase C delta (PKC delta) is a 78 kDa member of the novel group (nPKCs: sensitive to diacylglycerol, phosphatidylserine, and phorbol esters) of the PKC family of serine/threonine kinases that are involved in a wide range of physiological processes including mitogenesis, cell survival and transcriptional regulation. PKC delta is an ubiquitously expressed PKC isozyme that has been implicated in the regulation of multiple cellular processes including cell cycle progression and apoptosis. Autophosphorylation of serine 664 (serine 662 in mouse and rat) contributes to PKC delta activity. Serum dependent phosphorylation of serine 664 is mediated by mTOR pathway, is protected from dephosphorylation by phosphorylated threonine 505 in the activation loop, and is implicated in prolactin signaling.

# **Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:** 

~ 80 kDa

**Swiss-Prot:** 

Q05655

### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.


# **Specificity:**

p-PKC  $\delta$  (S645) polyclonal antibody detects endogenous levels of PKC  $\delta$  protein only when phosphorylated at Ser645.

#### **DATA:**



Western blot (WB) analysis of p-PKC δ (S645) pAb at 1:500 dilution Lane1:A549 whole cell lysate(20ug) Lane2:HCT116 whole cell lysate(20ug) Lane3:SGC7901 whole cell lysate(40ug) Lane4:The Uterus tissue lysate of Mouse(40ug) Lane5:The Uterus tissue lysate of Rat(40ug)



Immunohistochemistry (IHC) analyzes of p-PKC  $\delta$  (S645) pAb in paraffin-embedded human breast carcinoma tissue at 1:50.

## Note:

For research use only, not for use in diagnostic procedure.