

## GAPDH (1A6) monoclonal antibody

Catalog: BCP6605

Host: Mouse

Reactivity: Human, Mouse, Rat

### BackGround:

Glyceraldehyde 3 phosphate dehydrogenase (GAPDH) is well known as one of the key enzymes involved in glycolysis. As well as functioning as a glycolytic enzyme in cytoplasm, recent evidence suggests that mammalian GAPDH is also involved in a great number of intracellular processes such as membrane fusion, microtubule bundling, phosphotransferase activity, nuclear RNA export, DNA replication, and DNA repair. During the last decade a lot of data appeared concerning the role of GAPDH in different pathologies including prostate cancer progression, programmed neuronal cell death, age related neuronal diseases, such as Alzheimer's and Huntington's disease. GAPDH is expressed in all cells. It is constitutively expressed in almost all tissues at high levels.

### Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 36 kDa

### Swiss-Prot:

N/A

### Purification&Purity:

The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

ELISA: 1:50000~1:100000

WB: 1:10000

IHC:1:500~1:1000

IF:1:500~1:1000


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

GAPDH (1A6) mAb detects endogenous levels of GAPDH protein.

### DATA:



Western blot (WB) analysis of GAPDH (1A6) mAb at 1:20000/1:10000 dilution

Lane1:L02 whole cell lysate(40ug)

Lane2:A549 whole cell lysate(40ug)


Lane3:MG63 whole cell lysate(40ug)

Lane4:PC12 whole cell lysate(40ug)


Lane5:BV2 whole cell lysate(40ug)

Lane6:The Brain tissue lysate of Rat(40ug)

Lane7:The Brain tissue lysate of Mouse(40ug)



Immunohistochemistry analyzes of GAPDH (1A6) antibody in paraffin-embedded human liver tissue



IF image of BCP6605 stained A375 cells. The cells were 4% paraformaldehyde fixed (20 min) and then incubated in 10% normal goat serum for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody GAPDH (1A6) mAb #BCP6605(1:100) at 10 µg/ml overnight at +4 °C. The secondary antibody (GREEN) was Goat Anti-Mouse IgG (H+L) FITC used at a 1/400 dilution for 1h. Hoechst33342 was used to

stain the cell nuclei (blue).

For research use only, not for use in diagnostic procedure.

**Note:**